

PLURIPOLARITY OF GRAPHS OF ALGEBROID FUNCTIONS

ZAFAR S. IBRAGIMOV

Let $K \subset \mathbf{C}^n$ be an arbitrary compact set and let $f(z)$ be a continuous function on K . By $\rho_m(f, K)$ we denote the least deviation of $f(z)$ on K from the rational functions of degree less than or equal to m . That is,

$$\rho_m(f, K) = \inf_{r_m} \|f - r_m\|_K,$$

where $\| - \|_K$ is the uniform norm and the infimum is taken over all rational functions of the form

$$r_m(z) = \frac{\sum_{|\alpha| \leq m} a_\alpha z^\alpha}{\sum_{|\alpha| \leq m} b_\alpha z^\alpha}, \quad \text{where } \alpha = (\alpha_1, \alpha_2, \dots, \alpha_n) \text{ is a multiindex.}$$

As usual, we denote by $e_m(f, K)$ the least deviation of function $f(z)$ on K from the polynomials of degree less than or equal to m . Obviously,

$$\rho_m(f, K) \leq e_m(f, K) \quad \text{for each } m = 1, 2, \dots$$

In ([1], [2]) Gonchar proved that if $K = [a, b] \subset \mathbf{R} \subset \mathbf{C}$, then the class of functions

$$R([a, b]) = \{f \in C[a, b]: \varliminf_{m \rightarrow \infty} \sqrt[m]{\rho_m(f, K)} < 1\}$$

possesses one of the important properties of the class of analytic functions. Namely, if

$$\varliminf_{m \rightarrow \infty} \sqrt[m]{\rho_m(f, K)} < 1\}$$

and if $f(x) = 0$ on a set $E \subset [a, b]$ of positive logarithmic capacity, then $f(x) \equiv 0$ on $[a, b]$ (see also [5]).

By analogy with the class

$$B(K) = \{f \in C(K): \varliminf_{m \rightarrow \infty} \sqrt[m]{e_m(f, K)} < 1\},$$

Date: May 3, 2010.

2000 Mathematics Subject Classification. Primary 32U.

which is called the class of quasianalytic functions of Bernstein ([3],[4],[6]), we call

$$R(K) = \{f \in C(K) : \varliminf_{m \rightarrow \infty} \sqrt[m]{\rho_m(f, K)} < 1\}$$

the class of quasianalytic functions of Gonchar.

The classes $B(K)$ and $R(K)$ are not linear spaces; the sum of two quasianalytic functions are not, in general, quasianalytic (see [4]). We consider the following subclass $R_0(K)$ of the class $R(K)$:

$$R_0(K) = \{f \in C(K) : \overline{\lim}_{m \rightarrow \infty} \sqrt[m]{\rho_m(f, K)} < 1\}$$

It is not hard to see that, if f_1 and f_2 belong to $R_0(K)$, then so are $c_1 f_1 + c_2 f_2$ and $f_1 \cdot f_2$, where c_1 and c_2 are arbitrary complex numbers.

In ([7]) K. Diederich and J.E. Fornass constructed an example of a smooth (infinitely differentiable) function, whose graph is not pluripolar in $\mathbf{C}^2$. Recently, D. Coman, N. Levenberg and E.A. Poletskiy have proved, that if $f \in B([a, b])$, then its graph Γ_f is pluripolar in $\mathbf{C}^2$.

In ([8]) A. Edigarian studied the following analogue of a theorem of N. Shcherbina [9]. Let D be a domain in $\mathbf{C}^n$ and let $\Gamma \subset D \times \mathbf{C}$ be a graph of some algebroid function, i.e.,

$$\Gamma = \left\{ (z, w) \in D \times \mathbf{C} : w^k + a_1(z)w^{k-1} + \cdots + a_k(z) = 0 \right\},$$

where $a_1(z), a_2(z), \dots, a_k(z)$ are continuous functions on D . Then Γ is pluripolar in $\mathbf{C}^{n+1}$ if and only if the functions $a_1(z), a_2(z), \dots, a_k(z)$ are holomorphic in D . In this paper we prove a similar theorem on pluripolarity of graphs of algebroid functions in the class of quasianalytic functions.

Theorem 0.1. *Let $[a, b] \subset \mathbf{R} \subset \mathbf{C}$ and let $\Gamma \subset \mathbf{C}^2$ be a graph of some algebroid function, i.e.*

$$\Gamma = \left\{ (z, w) \in D \times \mathbf{C} : w^k + a_1(z)w^{k-1} + \cdots + a_k(z) = 0 \right\},$$

where $a_l \in R_0([a, b])$, $l = 1, 2, \dots, k$. Then Γ is pluripolar in $\mathbf{C}^2$

Proof. We consider the following function on $[a, b] \times \mathbf{C}$

$$f(z, w) = w^k + a_1(z)w^{k-1} + \cdots + a_k(z).$$

Since $a_l \in R_0([a, b])$ for $l = 1, 2, \dots, k$, there exist a sequence of rational functions $r_1^j(z), r_2^j(z), \dots, r_m^j(z), \dots$ such that

$$\sqrt[m]{\rho_m(a_j, [a, b])} = \sqrt[m]{\|a_j(z) - r_m^j(z)\|_{[a, b]}} \leq \delta_j < 1.$$

The function $f(z, w)$ is quasianalytic in the sense of Gonchar on the compact set $[a, b] \times \{|w| \leq h\} \subset \mathbf{C}^{k+1}$, where h is an arbitrary positive number. Indeed,

$$\begin{aligned} \rho_{m+k}(f, [a, b] \times \{|w| \leq h\}) &\leq \|f(z, w) - w^k - \sum_{j=1}^k r_m^j(z) w^{k-j}\|_{[a, b] \times \{|w| \leq h\}} \\ &\leq \sum_{j=1}^k \|a_j(z) - r_m^j(z)\|_{[a, b]} \cdot h^j \leq \sum_{j=1}^k h^j \delta_j^{m+k} \leq k \max\{h, h^k\} \delta^{m+k}, \end{aligned}$$

where $\delta = \max\{\delta_j : j = 1, 2, \dots, k\}$. It follows that

$$\overline{\lim}_{m \rightarrow \infty} \rho_{m+k}^{1/(m+k)}(f, [a, b] \times \{|w| \leq h\}) \leq \delta < 1.$$

Consequently, the graph

$$\Gamma_f = \{(z, w, f(z)) : (z, w) \in [a, b] \times \mathbf{C}\}$$

of the function $f(z, w)$ is pluripolar in $\mathbf{C}^3$. Now we consider sections

$$\Gamma_f(\lambda) = \{(z, w, f(z, w)) : f(z, w) = \lambda\}.$$

For each λ , the section $\Gamma_f(\lambda)$ is pluripolar in $\mathbf{C}^2$. (Indeed, if the graph $\Gamma_f(\lambda)$ is nonpluripolar for some $\lambda \in \mathbf{C}$, then according to the uniqueness property of quasianalytic functions, the function $f(z, w)$ is identically equal to λ , which contradicts the definition of $f(z, w)$). In particular, we obtain the pluripolarity of the graph

$$\Gamma = \{(z, w) \in D \times \mathbf{C} : w^k + a_1(z)w^{k-1} + \dots + a_k(z) = 0\}$$

of algebroid functions. The proof of the theorem is complete. $\square$

REFERENCES

- [1] A.A. Gonchar, *On best approximations by rational functions*, Dokl. Akad. Nauk SSSR, 100 (1955), 205–208, (Russian).
- [2] A.A. Gonchar, *Quasianalytic class of functions, connected with best approximated by rational functions*, Izv. of the Acad. of Sci. Armenia SSR. 1971, VI . No. 2-3, 148–159.
- [3] A.F. Timan, *Theory of approximation of functions of a real variable*, Pergaon Press, Macmillian, New York, 1963.

- [4] W. Plesniak, *Quasianalytic functions of several complex variables*, Zeszyty Nauk. Uniw. Jagiell. 15 (1971), pp. 135–145.
- [5] W. Plesniak, *Characterization of quasianalytic functions of several variables by means of rational approximation*, Ibid. 27 (1973), pp. 149–157.
- [6] D. Coman, N. Levenberg, E.A. Poletskiy, *Quasianalyticity and Pluripolarity*, J. Amer. Math. Soc. 18 (2005) No. 2, pp. 9–16.
- [7] K. Diederich, J.E. Fornass, *A smooth curve in which is not a pluripolar set*, Duke Math. J. 49 (1982), pp. 931–936.
- [8] A. Edigarian, *Graphs of multifunctions*, Math. Z., 250 (2005) 145–147.
- [9] N. Shcherbina, *Pluripolar graphs are holomorphic*, Acta Math., 194 (2005), pp. 203–216.

DEPARTMENT OF MATHEMATICS, URGENCH STATE UNIVERSITY, URGENCH 220100, UZBEK-
ISTAN

E-mail address: z.ibragim@gmail.com